



DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

mTORC1 stimulates nucleotide synthesis through both transcriptional and post-translational mechanisms

The Harvard community has made this article openly available.
[Please share](#) how this access benefits you. Your story matters.

Citation	Ben-Sahra, Issam, Stephane Ricoult, Jessica Howell, John Asara, and Brendan Manning. 2014. "mTORC1 stimulates nucleotide synthesis through both transcriptional and post-translational mechanisms." <i>Cancer & Metabolism</i> 2 (Suppl 1): P6. doi:10.1186/2049-3002-2-S1-P6. http://dx.doi.org/10.1186/2049-3002-2-S1-P6 .
Published Version	doi:10.1186/2049-3002-2-S1-P6
Accessed	February 16, 2015 3:27:54 PM EST
Citable Link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12717406
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

(Article begins on next page)

POSTER PRESENTATION

Open Access

mTORC1 stimulates nucleotide synthesis through both transcriptional and post-translational mechanisms

Issam Ben-Sahra^{1*}, Stephane Ricoult¹, Jessica Howell¹, John Asara², Brendan Manning¹

From Metabolism, Diet and Disease 2014: Cancer and metabolism
Washington DC, USA. 28-30 May 2014

Background

Cellular growth signals stimulate anabolic processes. The mechanistic target of rapamycin (mTOR), as part of mTORC1, is a protein kinase that senses growth signals to regulate anabolic growth and proliferation. mTORC1 stimulates protein synthesis through effects on mRNA translation and ribosome biogenesis [1]. mTORC1 signaling also promotes *de novo* lipid and sterol synthesis through the activation of the sterol-response element-binding protein (SREBP) transcription factors, which stimulate the expression of the enzymes driving this biosynthetic process [2].

Material and Methods

TSC2^{+/+} MEFs, *TSC2*^{-/-} MEFs, MCF10A expressing pBabe-empty vector or PI3KCA^{H1047R}, HeLa cells, U87MG cell line were used in this study. To determine the relative levels of intracellular metabolites, extracts were prepared and analyzed by LC/MS/MS [3]. Regarding the U-¹⁴C-aspartate, U-¹⁴C-glycine incorporation into RNA and DNA, cells were serum starved for 15 hours and treated as indicated. Cells were harvested and RNA or DNA was isolated using Allprep DNA/RNA kits according to the manufacturer's instructions and quantified using a spectrophotometer. For statistical analysis a two-tailed Student's t-test was performed for all pairwise comparisons (*n*=3).

Results

We find that activation of mTORC1 leads to the acute stimulation of metabolic flux through the *de novo* pyrimidine synthesis pathway [4]. We recently found that

mTORC1 stimulates the *de novo* purine synthesis pathway. In contrast with pyrimidine synthesis, the regulation of the purine synthesis by mTORC1 signaling occurs through long-term mechanism. Indeed, we found that mTORC1 regulates the *de novo* purine synthesis pathway through the transcription factor SREBP.

Conclusion

These findings demonstrate that growth signaling through mTORC1 promotes the production of new nucleotides to facilitate an increased demand for RNA and DNA. mTOR appears to be a central regulator of *de novo* nucleotide synthesis. Therefore, nucleotide synthesis joins protein and lipid synthesis as major anabolic processes stimulated by mTORC1 signaling.

Authors' details

¹Harvard School of Public Health, Department of Genetics and Complex Diseases, Boston, MA, USA. ²Division of Signal Transduction, Beth Israel Deaconess Medical Center and Department of Medicine, Harvard Medical School, Boston, MA, USA.

Published: 28 May 2014

References

1. Laplante M, Sabatini DM: mTOR signaling in growth control and disease. *Cell* 2012, **149**:274.
2. Düvel K, et al: Activation of a metabolic gene regulatory network downstream of mTOR complex 1. *Mol Cell* 2010, **39**:171.
3. Yuan M, et al: A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue. *Nat Protoc* 2012, **7**:872.
4. Ben-Sahra I, et al: Stimulation of *de novo* pyrimidine synthesis by growth signaling through mTOR and S6K1. *Science* 2013, **339**:1323-1328.

doi:10.1186/2049-3002-2-S1-P6

Cite this article as: Ben-Sahra et al.: mTORC1 stimulates nucleotide synthesis through both transcriptional and post-translational mechanisms. *Cancer & Metabolism* 2014 **2**(Suppl 1):P6.

¹Harvard School of Public Health, Department of Genetics and Complex Diseases, Boston, MA, USA

Full list of author information is available at the end of the article